Contents:

Volume 132, Numbers 1 & 2, January 1996

Mini-review

1 **Chaperonin-mediated folding of actin and tubulin.**
S.A. Lewis, G. Tian, I.E. Vainberg, and N.J. Cowan

Regular Articles

5 **Assembly of the nuclear pore: Biochemically distinct steps revealed with NEM, GTPγS, and BAPTA.**
C. Macaulay and D.J. Forbes

21 **A novel class of clathrin-coated vesicles budding from endosomes.**
W. Stoorvogel, V. Oorschot, and H.J. Geuze

35 **Endocytosis of GPI-linked membrane folate receptor-α.**
S. Rijnboutt, G. Jansen, G. Posthuma, J.B. Hynes, J.H. Schornagel, and G.J. Strous

49 **Phagosome–lysosome fusion is a calcium-independent event in macrophages.**
S. Zimmerli, M. Majeed, M. Gustavsson, O. Stendahl, D.A. Sanan, and J.D. Ernst

63 **A new chloroplast protein import intermediate reveals distinct translocation machineries in the two envelope membranes: Energetics and mechanistic implications.**
S.V. Scott and S.M. Theg

77 **Identification and localization of an actin-binding motif that is unique to the epsilon isoform of protein kinase C and participates in the regulation of synaptic function.**
R. Prekeris, M.W. Mayhew, J.B. Cooper, and D.M. Terrian

91 **Involvement of fyn tyrosine kinase in progression of cytokinesis of B lymphocyte progenitor.**
M. Yasunaga, T. Yagi, N. Hanzawa, M. Yasuda, Y. Yamanashi, T. Yamamoto, S. Aizawa, Y. Miyauchi, and S.-I. Nishikawa

101 ***Dictyostelium* myosin heavy chain kinase A regulates myosin localization during growth and development.**
M.F. Kolman, L.M. Futey, and T.T. Egelhoff

111 **The *Saccharomyces cerevisiae* spindle pole body duplication gene *MPS1* is part of a mitotic checkpoint.**
E. Weiss and M. Winey

125 **Initial triggering of M-phase in starfish oocytes: A possible novel component of maturation-promoting factor besides cdc2 kinase.**
E. Okumura, T. Sekiai, S.-I. Hisanaga, K. Tachibana, and T. Kishimoto

137 ***BED1*, a gene encoding a galactosyltransferase homologue, is required for polarized growth and efficient bud emergence in *Saccharomyces cerevisiae*.**
G. Mondésert and S.I. Reed

153 **A nontetrameric species is the major soluble form of keratin in *Xenopus* oocytes and rabbit reticulocyte lysates.**
J.B. Bachant and M.W. Klymkowsky

167 **Vamp/synaptobrevin isoforms 1 and 2 are widely and differentially expressed in nonneuronal tissues.**
O. Rossetto, L. Gorza, G. Schiavo, N. Schiavo, R.H. Scheller, and C. Montecucco

181 **Lysophosphatidic acid and bFGF control different modes in proliferating myoblasts.**
S. Yoshida, A. Fujisawa-Sehara, T. Taki, K. Arai, and Y. Nabeshima

195 **Increased expression of TGF-β2 in osteoblasts results in an osteoporosis-like phenotype.**
A. Erlebacher and R. Derynck

211 **β1D integrin displaces the β1A isoform in striated muscles: Localization at junctional structures and signaling potential in nonmuscle cells.**
A.M. Belkin, N.I. Zhidkova, F. Balzac, F. Altruda, D. Tomatis, A. Maier, G. Tarone, V.E. Koteliansky, and K. Burridge

227 **β1 integrin–dependent and –independent polymerization of fibronectin.**
K. Wennerberg, L. Lohikangas, D. Gullberg, M. Pfaff, S. Johansson, and R. Fässler

239 **Extracellular matrix alters PDGF regulation of fibroblast integrins.**
J. Xu and R.A.F. Clark

Volume 132, Number 3, February 1996

Mini-Review

255 **Roles of molecular chaperones in protein degradation.**
S.A. Hayes and J.F. Dice

Regular Articles

259 **XIST RNA paints the inactive X chromosome at interphase: Evidence for a novel RNA involved in nuclear/chromosome structure.**
C.M. Clemson, J.A. McNeil, H.F. Willard, and J.B. Lawrence

277 **Biochemical requirements for the targeting and fusion of ER-derived transport vesicles with purified yeast Golgi membranes.**
V.V. Lupashin, S. Hamamoto, and R.W. Schekman

291 **Assembly of ER-associated protein degradation in vitro: Dependence on cytosol, calnexin, and ATP.**
A.A. McCracken and J.L. Brodsky

299 **Role of CD3γ in T cell receptor assembly.**
J. Dietrich, A. Neisig, X. Hou, A.-M.K. Wegener, M. Gajhede, and C. Geisler

311 **Import of a DHFR hybrid protein into glycosomes in vivo is not inhibited by the folate-analogue aminopterin.**
T. Häusler, Y.-D. Stierhof, E. Wirtz, and C. Clayton

325 **Peb1p (Pas7p) is an intraperoxisomal receptor for the NH_2-terminal, type 2, peroxisomal targeting sequence of thiolase: Peb1p itself is targeted to peroxisomes by an NH_2-terminal peptide.**
J.W. Zhang and P.B. Lazarow

335 **Overexpression of cofilin stimulates bundling of actin filaments, membrane ruffling, and cell movement in *Dictyostelium*.**
H. Aizawa, K. Sutoh, and I. Yahara

345 **Removal of MAP4 from microtubules in vivo produces no observable phenotype at the cellular level.**
X.M. Wang, J.G. Peloquin, Y. Zhai, J.C. Bulinski, and G.G. Borisy

359 ***PF16* encodes a protein with armadillo repeats and localizes to a single microtubule of the central apparatus in *Chlamydomonas* flagella.**
E.F. Smith and P.A. Lefebvre

371 **Sequence and submolecular localization of the 115-kD accessory subunit of the heterotrimeric kinesin-II ($KRP_{85/95}$) complex.**
K.P. Wedaman, D.W. Meyer, D.J. Rashid, D.G. Cole, and J.M. Scholey

381 **Onset of re-epithelialization after skin injury correlates with a reorganization of keratin filaments in wound edge keratinocytes: Defining a potential role for keratin 16.**
R.D. Paladini, K. Takahashi, N.S. Bravo, and P.A. Coulombe

399 **Identification of a developmentally regulated septin and involvement of the septins in spore formation in *Saccharomyces cerevisiae*.**
H. Fares, L. Goetsch, and J.R. Pringle

413 **Mitotic mechanisms in Alzheimer's disease?**
I. Vincent, M. Rosado, and P. Davies

427 **TrkA receptor ectodomain cleavage generates a tyrosine-phosphorylated cell-associated fragment.**
N. Cabrera, E. Díaz-Rodríguez, E. Becker, D. Martín-Zanca, and A. Pandiella

437 **CD47 mediates post-adhesive events required for neutrophil migration across polarized intestinal epithelia.**
C.A. Parkos, S.P. Colgan, T. Liang, A. Nusrat, A.E. Bacarra, D.K. Carnes, and J.L. Madara

451 **Catenins and zonulus occulens-1 form a complex during early stages in the assembly of tight junctions.**
A.K. Rajasekaran, M. Hojo, T. Huima, and E. Rodriguez-Boulan

465 **Visualization of CD2 interaction with LFA-3 and determination of the two-dimensional dissociation constant for adhesion receptors in a contact area.**
M.L. Dustin, L.M. Ferguson, P.-Y. Chan, T.A. Springer, and D.E. Golan

475 **Human neural cell adhesion molecule L1 and rat homologue NILE are ligands for integrin $\alpha_v\beta_3$.**
A.M.P. Montgomery, J.C. Becker, C.-H. Siu, V.P. Lemmon, D.A. Cheresh, J.D. Pancook, X. Zhao, and R.A. Reisfeld

487 **Heparan sulfate expression in polarized epithelial cells: The apical sorting of glypican (GPI-anchored proteoglycan) is inversely related to its heparan sulfate content.**
G. Mertens, B. van der Schueren, H. van den Berghe, and G. David

Volume 132, Number 4, February 1996

499 **ER membrane protein complex required for nuclear fusion.**
D.T.W. Ng and P. Walter

511 **Activation of transcription factor NF-κB by the adenovirus E3/19K protein requires its ER retention.**
H.L. Pahl, M. Sester, H.-G. Burgert, and P.A. Baeuerle

523 **The AP-1 adaptor complex binds to immature secretory granules from PC12 cells, and is regulated by ADP-ribosylation factor.**
A.S. Dittié, N. Hajibagheri, and S.A. Tooze

537 **Endocytosis of VAMP is facilitated by a synaptic vesicle targeting signal.**
E. Grote and R.B. Kelly

549 **The *PALI* gene product is a peroxisomal ATP-binding cassette transporter in the yeast *Saccharomyces cerevisiae*.**
E.E. Swartzman, M.N. Viswanathan, and J. Thorner

565 **The targeting of Lamp1 to lysosomes is dependent on the spacing of its cytoplasmic tail tyrosine sorting motif relative to the membrane.**
J. Rohrer, A. Schweizer, D. Russell, and S. Kornfeld

577 **Cysteine^{34} of the cytoplasmic tail of the cation-dependent mannose 6-phosphate receptor is reversibly palmitoylated and required for normal trafficking and lysosomal enzyme sorting.**
A. Schweizer, S. Kornfeld, and J. Rohrer

585 **Different fates of phagocytosed particles after delivery into macrophage lysosomes.**
Y.-K. Oh and J.A. Swanson

595 **Potential sites of PI-3 kinase function in the endocytic pathway revealed by the PI-3 kinase inhibitor, wortmannin.**
H. Shpetner, M. Joly, D. Hartley, and S. Corvera

607 **Calcium release in HSY cells conforms to a steady-state mechanism involving regulation of the inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[Ca^{2+}]$.**
A. Tanimura and R.J. Turner

617 **Molecular characterization of the 50-kD subunit of dynactin reveals function for the complex in chromosome alignment and spindle organization during mitosis.**
C.J. Echeverri, B.M. Paschal, K.T. Vaughan, and R.B. Vallee

635 **Detection of protein kinase activity specifically activated at metaphase-anaphase transition.**
M. Sekimata, K. Tsujimura, J. Tanaka, Y. Takeuchi, N. Inagaki, and M. Inagaki

643 **Filensin and phakinin form a novel type of beaded intermediate filaments and coassemble de novo in cultured cells.**
G. Goulielmos, F. Gounari, S. Remington, S. Müller, M. Häner, U. Aebi, and S.D. Georgatos

657 **Myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis.**
V. Andrés and K. Walsh

667 **Selective stabilization of tau in axons and microtubule-associated protein 2C in cell bodies and dendrites contributes to polarized localization of cytoskeletal proteins in mature neurons.**
N. Hirokawa, T. Funakoshi, R. Sato-Harada, and Y. Kanai

681 **Tenascin-C contains distinct adhesive, anti-adhesive, and neurite outgrowth promoting sites for neurons.**
B. Götz, A. Scholze, A. Clement, A. Joester, K. Schütte, F. Wigger, R. Frank, E. Spiess, P. Ekblom, and A. Faissner

701 **Human multidrug resistance 3–P-glycoprotein expression in transgenic mice induces lens membrane alterations leading to cataract.**
I. Dunia, J.J.M. Smit, M.A. van der Valk, H. Bloemendal, P. Borst, and E.L. Benedetti

717 **MDC9, a widely expressed cellular disintegrin containing cytoplasmic SH3 ligand domains.**
G. Weskamp, J. Krätzschmar, M.S. Reid, and C.P. Blobel

727 **Anchorage mediated by integrin α6β4 to laminin 5 (epiligrin) regulates tyrosine phosphorylation of a membrane-associated 80-kD protein.**
Y. Xia, S.G. Gil, and W.G. Carter

741 **Stimulation of β_1 integrins on fibroblasts induces PDGF independent tyrosine phosphorylation of PDGF β-receptors.**
C. Sundberg and K. Rubin

753 **ADDITIONS AND CORRECTIONS**

Volume 132, Number 5, March 1996

755 **Assembly of the ER to Golgi SNARE complex requires Uso1p.**
S.K. Sapperstein, V.V. Lupashin, H.D. Schmitt, and M.G. Waters

769 **The biogenesis of the MHC class II compartment in human I-Cell disease B lymphoblasts.**
J.N. Glickman, P.A. Morton, J.W. Slot, S. Kornfeld, and H.J. Geuze

787 **Thioredoxin is required for vacuole inheritance in *Saccharomyces cerevisiae*.**
Z. Xu and W. Wickner

795 **An internalization signal in the simian immunodeficiency virus transmembrane protein cytoplasmic domain modulates expression of envelope glycoproteins on the cell surface.**
M.M. Sauter, A. Pelchen-Matthews, R. Bron, M. Marsh, C.C. LaBranche, P.J. Vance, J. Romano, B.S. Haggarty, T.K. Hart, W.M.F. Lee, and J.A. Hoxie

813 **Sorting of newly synthesized galactosphingolipids to the two surface domains of epithelial cells.**
P. van der Bijl, M. Lopes-Cardozo, and G. van Meer

823 **Two-dimensional crystals of photosystem II: Biochemical characterization, cryoelectron microscopy and localization of the D1 and cytochrome b559 polypeptides.**
K.M. Marr, D.N. Mastronarde, and M.K. Lyon

835 **The *Caenorhabditis elegans* gene *unc-89*, required for muscle M-line assembly, encodes a giant modular protein composed of Ig and signal transduction domains.**
G.M. Benian, T.L. Tinley, X. Tang, and M. Borodovsky

849 **Expression of the *SM22α* promoter in transgenic mice provides evidence for distinct transcriptional regulatory programs in vascular and visceral smooth muscle cells.**
L. Li, J.M. Miano, B. Mercer, and E.N. Olson

861 **Movement of cortical actin patches in yeast.**
J.A. Waddle, T.S. Karpova, R.H. Waterston, and J.A. Cooper

871 ***Xenopus laevis* actin-depolymerizing factor/cofilin: A phosphorylation-regulated protein essential for development.**
H. Abe, T. Obinata, L.S. Minamide, and J.R. Bamburg

887 **Spc42p: A phosphorylated component of the *S. cerevisiae* spindle pole body (SPB) with an essential function during SPB duplication.**
A.D. Donaldson and J.V. Kilmartin

903 **The 110-kD spindle pole body component of *Saccharomyces cerevisiae* is a phosphoprotein that is modified in a cell cycle–dependent manner.**
D.B. Friedman, H.A. Sundberg, E.Y. Huang, and T.N. Davis

915 **A cell cycle–associated change in Ca^{2+} releasing activity leads to the generation of Ca^{2+} transients in mouse embryos during the first mitotic division.**
T. Kono, K.T. Jones, A. Bos-Mikich, D.G. Whittingham, and J. Carroll

925 **Gene targeting at the mouse cytokeratin 10 locus: Severe skin fragility and changes of cytokeratin expression in the epidermis.**
R.M. Porter, S. Leitgeb, D.W. Melton, O. Swensson, R.A.J. Eady, and T.M. Magin

937 **Agrin-induced acetylcholine receptor clustering in mammalian muscle requires tyrosine phosphorylation.**
M. Ferns, M. Deiner, and Z. Hall

945 **The neurotrophin receptor, gp75, forms a complex with the receptor tyrosine kinase TrkA.**
A.H. Ross, M.-C. Daou, C.A. McKinnon, P.J. Condon, M.B. Lachyankar, R.M. Stephens, D.R. Kaplan, and D.E. Wolf

955 **Bcl-2 inhibits retinoic acid–induced apoptosis during the neural differentiation of embryonal stem cells.**
H. Okazawa, J. Shimizu, M. Kamei, I. Imafuku, H. Hamada, and I. Kanazawa

969 **Contractile activity regulates isoform expression and polysialylation of NCAM in cultured myotubes: Involvement of Ca^{2+} and protein kinase C.**
V.F. Rafuse and L. Landmesser

Volume 132, Number 6, March 1996

985 **Localization of a yeast early Golgi mannosyltransferase, Och1p, involves retrograde transport.**
S.L. Harris and M.G. Waters

999 **Identification of a cytoplasm to vacuole targeting determinant in aminopeptidase I.**
M.N. Oda, S.V. Scott, A. Hefner-Gravink, A.D. Caffarelli, and D.J. Klionsky

1011 **Multivesicular endosomes containing internalized EGF–EGF receptor complexes mature and then fuse directly with lysosomes.**
C.E. Futter, A. Pearse, L.J. Hewlett, and C.R. Hopkins

1025 **Endocytosis of activated receptors and clathrin-coated pit formation: Deciphering the chicken or egg relationship.**
F. Santini and J.H. Keen

1037 **Na^+/H^+ exchange activity during phagocytosis in human neutrophils: Role of Fcγ receptors and tyrosine kinases.**
T. Fukushima, T.K. Waddell, S. Grinstein, G.G. Goss, J. Orlowski, and G.P. Downey

1053 **Biochemical kinetic characterization of the *Acanthamoeba* myosin-I ATPase.**
E.M. Ostap and T.D. Pollard

1061 **Developmental genetic analysis of troponin T mutations in striated and nonstriated muscle cells of *Caenorhabditis elegans*.**
C.D. Myers, P.-Y. Goh, T. StC. Allen, E.A. Bucher, and T. Bogaert

1079 **Identification, localization, and functional implications of an abundant nematode annexin.**
C.E. Creutz, S.L. Snyder, S.N. Daigle, and J. Redick

1093 **The force for poleward chromosome motion in *Haemanthus* cells acts along the length of the chromosome during metaphase but only at the kinetochore during anaphase.**
A. Khodjakov, R.W. Cole, A.S. Bajer, and C.L. Rieder

1105 **Binding to cadherins antagonizes the signaling activity of β-catenin during axis formation in *Xenopus*.**
F. Fagotto, N. Funayama, U. Glück, and B.M. Gumbiner

1115 **The estrogen-dependent c-JunER protein causes a reversible loss of mammary epithelial cell polarity involving a destabilization of adherens junctions.**
I. Fialka, H. Schwarz, E. Reichmann, M. Oft, M. Busslinger, and H. Beug

1133 **Population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, EGF and TGFα in a chemically defined (HGM) medium.**
G.D. Block, J. Locker, W.C. Bowen, B.E. Petersen, S. Katyal, S.C. Strom, T. Riley, T.A. Howard, and G.K. Michalopoulos

1151 **Differentially expressed fibroblast growth factors regulate skeletal muscle development through autocrine and paracrine mechanisms.**
K. Hannon, A.J. Kudla, M.J. McAvoy, K.L. Clase, and B.B. Olwin

1161 **Cloning of mouse integrin α_v cDNA and role of the α_v-related matrix receptors in metanephric development.**
J. Wada, A. Kumar, Z. Liu, E. Ruoslahti, L. Reichardt, J. Marvaldi, and Y.S. Kanwar

1177 **Cooperative effect of TNFα, bFGF, and VEGF on the formation of tubular structures of human microvascular endothelial cells in a fibrin matrix. Role of urokinase activity.**
P. Koolwijk, M.G.M. van Erck, W.J.A. de Vree, M.A. Vermeer, H.A. Weich, R. Hanemaaijer, and V.W.M. van Hinsbergh

1189 **Human amnion contains a novel laminin variant, laminin 7, which like laminin 6, covalently associates with laminin 5 to promote stable epithelial–stromal attachment.**
M.-F. Champliaud, G.P. Lunstrum, P. Rousselle, T. Nishiyama, D.R. Keene, and R.E. Burgeson

1199 **Glycosylation of CD44 is implicated in CD44-mediated cell adhesion to hyaluronan.**
A. Bartolazzi, A. Nocks, A. Aruffo, F. Spring, and I. Stamenkovic

1209 **Syndecan-1 mediates cell spreading in transfected human lymphoblastoid (Raji) cells.**
C.S. Lebakken and A.C. Rapraeger

1223 **AUTHOR INDEX FOR VOLUME 132**

1231 **SUBJECT INDEX FOR VOLUME 132**